

Biologically Viable Methods for Food Preservation: A Review

Subha Ganguly

AICRP on Post Harvest Technology (ICAR), Department of Fish Processing Technology,
Faculty of Fishery Sciences, West Bengal University of Animal and Fishery Sciences, 5, Budherhat Road,
P.O. Panchasayar, Chakgaria, Kolkata 700 094, WB, India
E-mail: ganguly38@gmail.com

ABSTRACT

Preservation of food is aimed towards prevention and control of growth of spoilage microorganisms and bacteria including fungi and yeasts and also reducing the rancidity due to oxidation of lipids. Food preservation is also targeted towards reduction of discoloration of food products by enzymatic browning which occurs during preparation of processed food products. The present article discusses on the biologically and ecologically viable methods for food preservation which is gaining importance in food processing industries.

Key words: Biological, Food, Preservation

Received 26/04/2013 Accepted 01/05/2013

© 2013 Academy for Environment and Life Sciences, India

INTRODUCTION

Biopreservation is the method used for food preservation by using natural antimicrobials and microbiota thereby increasing the storage life of food [1]. The beneficial products formed due to fermentation by the bacteria are used in this process to reduce the rate of food spoilage and to render the food free from pathogenic microorganisms and metabolites [2, 9]. This process is gradually increasing in popularity for its ecologically benign approach [1].

Lactic acid bacteria are important in this aspect as natural biopreservatives which possess antagonistic properties against the spoilage bacteria and pathogens [12]. The metabolites of LAB include acidic components like acetic acid, lactic acid, hydrogen peroxide and bacteriocins which are peptide in nature [12]. When the LAB competes for nutrients then they produce these metabolite components and the antimicrobial product, nisin which acts as a promising preservative for food [3, 4, 11].

The bacteriocins produced by LAB are also used in the Hurdle Technology. Using the bacteriocin producing LAB with other effective preservation techniques is effective in controlling the growth of spoilage microorganisms and inhibiting their generation, growth and bioactivity [1, 10, 12].

HURDLE TECHNOLOGY

This is a method of rendering the food to be free from contaminating and spoilage bacteria and pathogens by the combination of one or more methods. The pathogenic microorganisms have to pass through these individual approaches called "hurdles" for maintaining their activity in food products. Proper combination of hurdles will lead to destruction of the microbes and can prevent their further growth [5]. The proper combination of the hurdles ensures the microbial safety in food [8] thereby maintaining its nutritional and organoleptic parameters for consumer preference [6, 7].

The hurdles include the properties like processing at high temperature, storage at low temperature, lowering pH (increasing acidity), water activity (a_w) and/ or redox potential including the presence of biopreservatives or other preservative components in food products [13, 14]. The intensity of the hurdle is ascertained and controlled according to the type of spoilage microorganism(s) and regulated as per consumer safety and preference without sacrificing the quality and appearance of the final food product [5, 13, 14].

REFERENCES

1. Ananou, S., Maqueda, M., Martínez-Bueno, M., and Valdivia, E. (2007) Biopreservation, an ecological approach to improve the safety and shelf-life of foods. In: Méndez-Vilas, A. (Ed.) *Communicating Current Research and Educational Topics and Trends in Applied Microbiology*. Formatex. [ISBN 978-84-611-9423-0].

2. Yousef, A.E. and Carolyn Carlstrom, C. (2003) *Food microbiology: a laboratory manual*. Wiley, pp.226 [ISBN 978-0-471-39105-0].
3. FAO. Preservation techniques Fisheries and aquaculture department, Rome. Updated 27 May 2005. Retrieved 14 March 2011.
4. Alzamora, S.M., Tapia, M.S. and López-Malo, A. (2000) Minimally processed fruits and vegetables: fundamental aspects and applications. Springer, pp.266 [ISBN 978-0-8342-1672-3].
5. Alasalvar, C. (2010) *Seafood Quality, Safety and Health Applications*. John Wiley and Sons, pp.203. ISBN 978-1-4051-8070-2.
6. Leistner, I. (2000) Basic aspects of food preservation by hurdle technology. *International Journal of Food Microbiology*. **55**:181–186.
7. Leistner, L. (1995) Principles and applications of hurdle technology. In: Gould GW (Ed.) *New Methods of Food Preservation*, Springer, pp.1-21 [ISBN 978-0-8342-1341-8].
8. Ganguly, S. (2012) *Food Microbiology*. LAP LAMBERT Academic Publishing GmbH & Co. KG, Saarbrücken, Germany [ISBN 978-3-8484-8217-7].
9. Ganguly, S. (2012) *Food Processing Technology*. AV Akademikerverlag GmbH & Co. KG, Saarbrücken, Germany [ISBN 978-3-8383-8653-9].
10. L. De Vuyst and F. Leroy. (2007). Bacteriocins from lactic acid bacteria: production, purification, and food applications. *J. Mol. Microbiol. Biotechnol.* 13(4), 194-9.
11. Ingolf F. Nes, Dzung Bao Diep, Leiv Sigve Håvarstein, May Bente Brurberg, Vincent Eijsink and Helge Holo (1996) Biosynthesis of bacteriocins in lactic acid bacteria. *Antonie van Leeuwenhoek*. **70**(2-4), 113-128.
12. Cintas, L. M., Casaus, M. P., Herranz, C., Nes, I. F. and Hernández, P. E. (2001) Review: Bacteriocins of Lactic Acid Bacteria. doi: 10.1106/R8DE-P6HU-CLXP-5RYT *Food Science and Technology International*. 7(4), 281-305.
13. En Yang, Lihua Fan, Yueming Jiang, Craig Doucette and Sherry Fillmore (2012) Antimicrobial activity of bacteriocin-producing lactic acid bacteria isolated from cheeses and yogurts. *AMB Express*. 2, 48. doi:10.1186/2191-0855-2-48.
14. Pal, V., Jamuna, M. and Jeevaratnam, K. (2005) Isolation and characterization of bacteriocin producing lactic acid bacteria from a South Indian special dosa (Appam) batter. *Journal of Culture Collections*. 4, 53-60.